



Notes and Corrections on “Asymptotic Constancy Criteria of Solution of Linear Parabolic Volterra Difference Equations”

Z. Q. ZHANG

Department of Appl. Math., Hunan University

Changsha 410082, P.R. China

(Received May 1997; accepted July 1997)

Abstract—In this paper, we make some notes and corrections on “Asymptotic Constancy Criteria of Solutions of Linear Parabolic Volterra Difference Equations” [1].

Keywords—Parabolic Volterra difference equations, Neutral type, Asymptotic constancy, Von Neumann boundary conditions, Positive and negative coefficients.

In the recent paper, Shi *et al.* [1] studied the asymptotic constancy of solutions of linear parabolic Volterra difference equations of neutral type of form

$$\Delta_2 \left(u_{m,n} - \sum_{k=1}^{\infty} r_{k,n} u_{m,n-\alpha_k} \right) + \sum_{i=1}^{\infty} p_{i,n} u_{m,n-\beta_i} - \sum_{j=1}^{\infty} q_{j,n} u_{m,n-\gamma_j} = d \Delta_1^2 u_{m-1,n+1}, \quad (1)$$

$$\text{for } m = 1, \dots, M \text{ and } n \in z^+(n_0) = n_0, n_0 + 1, \dots,$$

with the homogeneous Von Neumann boundary conditions (NBC)

$$\Delta_1 u_{0,n} = \Delta_1 u_{M,n} = 0, \quad \text{for } n \in z^+(n_0), \quad (2)$$

and initial conditions (IC)

$$u_{m,l} = \theta_{m,l}, \quad \text{for } m = 1, \dots, M \text{ and } l \in \bar{z}(n_0) = \{\dots, n_0 - 1, n_0\}, \quad (3)$$

satisfies

$$\|\Theta\| = \sup \{|\Theta_{m,l}| \text{ for } m = 1, \dots, M \text{ and } l \in \bar{z}(n_0)\} < \infty, \quad (4)$$

where Δ_1 , Δ_1^2 , and Δ_2 are forward partial difference operators such that $\Delta_1 u_{m,n} = u_{m+1,n} - u_{m,n}$, $\Delta_1^2 u_{m,n} = \Delta_1(\Delta_1 u_{m,n})$, and $\Delta_2 u_{m,n} = u_{m,n+1} - u_{m,n}$ for $m = 1, \dots, M, n \in z = \{\dots, -1, 0, 1, \dots\}$. In the following, Shi *et al.* [1] assumed that $r_{k,n} \in z^+(1) \times z^+(n_0) \rightarrow R$, $p_{i,n} \in z^+(1) \times z^+(n_0) \rightarrow R^+$, $q_{j,n} \in z^+(1) \times z \rightarrow R^+$, $B_i \geq \gamma_i$ for $i \in z^+(1)$, $d \in R^+$ and that there exists a constant $c > 0$ such that $h_{i,n} \geq cq_{i,n-\beta_i+\gamma_i}$ for $i \in z^+(1)$ and sufficiently large n , where $h_{i,n} = p_{i,n} - q_{i,n-\beta_i+\gamma_i}$.

By constructing Liapunov sequences, Shi *et al.* [1] obtained the following asymptotic constancy theorem for equation (1).

THEOREM 1. (See [1, Theorem 2.1].) Assume that

$$\limsup_{n \rightarrow \infty} \left(\sum_{k=1}^{\infty} |r_{k,n+1}| + \sum_{j=1}^{\infty} \sum_{\mu=n+1-\beta_j+\gamma_j}^n q_{j,\mu} \right) = \delta < 1, \quad (c_1)$$

$$\begin{aligned} \limsup_{n \rightarrow \infty} \left[\delta \left(\frac{1 + \sum_{i=1}^{\infty} h_{i,n+1+\alpha_i+\beta_i}}{\sum_{i=1}^{\infty} h_{i,n+1+\beta_i}} \right) + \sum_{i=1}^{\infty} \sum_{\lambda=n+1-\beta_i}^{n+1} h_{i,\lambda+\beta_i} \right. \\ \left. + \frac{\sum_{i=1}^{\infty} (q_{i,n+1+\gamma_i} + h_{i,n+1+\beta_i}) \sum_{\lambda=n+2-\beta_i}^{n+1} h_{i,\lambda+2\beta_i}}{\sum_{i=1}^{\infty} h_{i,n+1-\beta_i}} \right] < 2, \end{aligned} \quad (c_2)$$

and that

$$\begin{aligned} \limsup_{n \rightarrow \infty} \left[\delta + \sum_{i=1}^{\infty} \sum_{\lambda=n+1-\beta_i}^n h_{i,\lambda+\beta_i} + \sum_{k=1}^{\infty} |r_{k,n+1+\alpha_k}| \right. \\ \left. + \sum_{i=1}^{\infty} \beta_i (q_{i,n+1+\gamma_i} + h_{i,n+1+\beta_i}) \right] < 2. \end{aligned} \quad (c_3)$$

Then every solution of IBVP (1)–(3) tends to a constant as $n \rightarrow \infty$.

One can point out in the proof of [1, Theorem 2.1], there are very serious errors such that the theorem does not hold. Now abstract a set of sentences in [1, pp. 74,75] as follows.

By Step II, we know

$$\sum_{m=1}^M \left(u_{m,n} - \sum_{k=1}^{\infty} r_{k,n} u_{m,n-\alpha_k} - \sum_{j=1}^{\infty} \sum_{\mu=n-\beta_j+\gamma_j}^{n-1} q_{j,\mu} u_{m,\mu-\gamma_j} \right) \rightarrow \sqrt{M\omega}, \quad \text{as } n \rightarrow \infty.$$

We will now prove that $\sum_{m=1}^M u_{m,n} \rightarrow \sqrt{M\omega}/(1-\delta)$ as $n \rightarrow \infty$. In the matter of fact, we let

$$\limsup_{n \rightarrow \infty} \sum_{m=1}^M u_{m,n} = \alpha \quad \text{and} \quad \liminf_{n \rightarrow \infty} \sum_{m=1}^M u_{m,n} = \beta.$$

Choose a subsequence $\{n_i\}$ of $\{n\}$ such that $\lim_{i \rightarrow \infty} \sum_{m=1}^M u_{m,n_i} = \alpha$. Let $\varepsilon > 0$ be sufficiently small such that

$$\begin{aligned} \sum_{m=1}^M u_{m,n_i} &= \sum_{m=1}^M \left(u_{m,n_i} - \sum_{k=1}^{\infty} r_{k,n_i} u_{m,n_i-\alpha_k} - \sum_{j=1}^{\infty} \sum_{\mu=n_i-\beta_j+\gamma_j}^{n_i-1} q_{j,\mu} u_{m,\mu-\gamma_j} \right) \\ &\quad + \sum_{m=1}^M \left(\sum_{k=1}^{\infty} r_{k,n_i} u_{m,n_i-\alpha_k} + \sum_{j=1}^{\infty} \sum_{\mu=n_i-\beta_j+\gamma_j}^{n_i-1} q_{j,\mu} u_{m,\mu-\gamma_j} \right) \\ &\leq \sqrt{M\omega} (\alpha + \varepsilon) \delta, \end{aligned} \quad (c_4)$$

$$(c_5)$$

and $\alpha \leq \sqrt{M\omega} + (\alpha + \varepsilon)\delta$. Letting $\varepsilon \rightarrow 0$, we obtain $\alpha \leq \sqrt{M\omega} + \alpha\delta$, i.e., $\alpha \leq \sqrt{M\omega}/(1-\delta)$. Similarly, we can show that

$$\beta \geq \frac{\sqrt{M\omega}}{1-\delta}. \quad (c_6)$$

Hence, $\alpha = \beta = \sqrt{M\omega}/(1-\delta)$.

One can see that since $\{u_{m,n}\}$ is assumed to be any solution of IBVP (1)–(3), the sign of α cannot be determined. Thus, after

$$\sum_{m=1}^M u_{m,n_i-\alpha_k} \quad \text{and} \quad \sum_{\mu=n_i-\beta_i+\gamma_i}^{n_i-1} u_{m,\mu-\gamma_i},$$

having been magnified into $(\alpha + \varepsilon)$, when

$$\sum_{k=1}^{\infty} |r_{k,n}| + \sum_{j=1}^{\infty} \sum_{\mu=n_i-\beta_i+\gamma_i}^{n_i-1} q_{j,\mu} u_{m,\mu-\gamma_i}$$

magnified into δ , it cannot be determined that whether (c_4) is magnified or reduced. Hence, (c_4) cannot follow (c_5) unless (c_1) is changed into

$$\lim_{n \rightarrow \infty} \left(\sum_{k=1}^{\infty} |r_{k,n+1}| + \sum_{j=1}^{\infty} \sum_{\mu=n+1-\beta_j+\gamma_j}^{n-1} q_{j,\mu} \right) = \delta < 1. \quad (5)$$

Similarly, since the sign of β cannot be determined, it cannot be defined whether (c_4) is magnified or reduced. Further, since $r_{k,n} \in Z^+(1) \times z^+(n_0) \rightarrow R$, from (c_4) , only get

$$\beta \geq \sqrt{M\omega} + \sum_{m=1}^M \left(\sum_{k=1}^{\infty} |r_{k,n}| u_{m,n_i-\alpha_k} - \sum_{j=1}^{\infty} \sum_{\mu=n_i-\beta_j+\gamma_j}^{n_i-1} q_{j,\mu} |u_{m,\mu-\gamma_j}| \right).$$

Thus, from (c_4) , one cannot get (c_6) . Hence, (c_4) cannot follow (c_6) unless (c_1) is changed into (c_5) and $r_{k,n} \in Z^+(1) \times Z^+(n_0) \rightarrow R$ is changed into $r_{k,n} \in z^+(1) \times z^+(n_1) \rightarrow R^+$.

The above mentioned errors can be avoided and the result presented in [1] remains a valid subject to the following corrections: (c_4) is changed into (c_5) and $r_{k,n} \in z^+(1) \times z^+(n_0) \rightarrow R$ is changed into $r_{k,n} \in z^+(1) \times z^+(n_1) \rightarrow R^+$.

REFERENCES

1. B. Shi *et al.*, Asymptotic constancy of solutions of linear parabolic Volterra difference equations, *Computers Math. Applic.* **32** (8), 65–77, (1996).
2. J.H. Shen and J.S. Yu, Asymptotic behavior of solutions of neutral differential equations with positive and negative coefficients, *J. Math. Anal. Appl.* **195**, 517–526, (1995).